

# useR Conference 2009

## *Impact Evaluation of Interventions on Child Health in Nepal*

Ron Bose PhD

Economist and Technical Officer

**3ie**

Rennes, France

July 7, 2009

# Diarrhea Prevalence in Nepal

1

**Table: 2001 Child Diarrhea Prevalence**

Response	Number	(%)
None	5,086	79
<b>Yes</b>	1,285	<b>20</b>
Total	6,415	100

*Source:* DHS 2001

2

**Table: 2006 Child Diarrhea Prevalence**

Response	Number	(%)
None	4,757	87
<b>Yes</b>	659	<b>12</b>
Total	5,457	100

*Source:* DHS 2006

# Access to Drinking Water

1

**Table: 2001 Water Source**

Source	Number	(%)
Piped Water	485	7
Public tap	1,825	26
Pvt. Well	135	2
Public Well	133	2
Tubewell	1,288	19
Public tubewell	1,177	17
Sprong/kuwa	1,267	18
River/lake/pond	166	2
Stone tap/dhara	58	1
Not resident	393	6
Total	6,929	100

*Source: DHS 2001*

2

**Table: 2006 Water Source**

Source	Number	(%)
Piped Water	513	9
Public tap	1,361	24
Pvt. well	25	0
Public well	140	2
Tubewell	2,044	35
Protected spring	144	2
Unprotected spring	640	11
River/dam/pond	376	7
Stone tap/dhara	205	4
Not dejure resident	318	5
Total	5,783	100

*Source: DHS 2006*

# Access to Sanitation

1

**Table: 2001 Toilet Facility**

Type	Number	(%)
Flush Toilet	511	7
Trad. Pit Toilet	971	14
Vent. Pit latrine	116	2
No facility	4,940	71
Not resident	393	6
<b>Total</b>	<b>6,931</b>	<b>100</b>

*Source: DHS 2001*

2

**Table: 2006 Toilet Facility**

Type	Number	(%)
Flush Toilet	1192	21
Trad Pit Toilet	909	15
Vent. Pit Latrine	48	1
No facility	3,250	56
Not dejure resident	318	5
<b>Total</b>	<b>5,782</b>	<b>100</b>

*Source: DHS 2006*

# Diarrhea Prevalence By Child Age in Months

1

Mean = 24.1 Months

Median = 21 Months

2

Mean = 23.13 Months

Median = 19 Months

# Diarrhea Prevalence: Access to "Improved Sanitation"

1

	Diarrhea	
	1	0
Imp. Toilet		
1	111	1131
0	548	3993

Source: DHS 2006

2

## Odds Ratio

$$\frac{\frac{P_1}{1-P_1}}{\frac{P_0}{1-P_0}} = 1.46$$

# Naive Comparison: Access to "Improved Sanitation"

I

**Table: Naive Comparison: Household Characteristics**

<b>Variable</b>	<b>Treatment</b>	<b>(Untreated)</b>
<i>Pipewtr. in house?</i>	23.2%	5%
<i>Rural</i>	52%	84%
<i>Head Hd has sec. or more ed.</i>	56%	30%
<i>House Floor= Cement</i>	29%	3%
<i>Richest Quintile</i>	54%	4%

*Source: DHS 2006*

# Rubin Neyman Causal Model

1

**Fundamental problem** with program evaluation is that it is physically impossible to observe counterfactual

2

Rubin (1974) gave us the model of identification of causal effects, which relies on the notion of a **synthetic counterfactual** for each observation. The model is based on work by Neyman (1923,1935) and Fisher (1918,1925); see also Tukey (1954), Wold (1956), Cochran (1965), Pearl (2000), and Rosenbaum (2002).

# Matching

- *Basic idea of matching is to compare outcome of treated and untreated individuals with similar  $x$ 's and then aggregating across  $x$ 's to get population average treatment effect. Advantage to regression approach is that it does not assume  $x$ 's linearly effect outcomes.*
- ***Propensity score matching (PSM)***  
 $\Delta^M = \frac{1}{N_T} \sum_{i \in (D=1)} [y_{1,i} - \sum_j w(i, j) y_{0,j}]$  *is to estimate the propensity score from the data, and then use that estimate to weight treatment effects for each propensity score accordingly to arrive at average treatment effect.*

# Comparison of Groups: Before versus After Matching

1

**Table: After Matching: Balanced Household Characteristics**

<b>Variable</b>	<b>Treatment</b>	<b>(Untreated)</b>
<i>Pipewtr. in house?</i>	23.2%	15%
<i>Rural</i>	53%	58%
<i>Head Hd has sec. or more ed.</i>	45%	41%
<i>House Floor= Cement</i>	30%	33%
<i>Richest Quintile</i>	52%	52%

*Source: DHS 2006*

# Impact Evaluation: Kernel Matching Results

1

Table: 2006 Results for Intervention on Diarrhea

Variable	Treatment	(Control)	$\Delta$	S.E.
Unmatched	0.091	0.122	-.032	(0.01)**
Matched	0.091	.143	-0.0524	(0.02)**

Note: "Treatment" = *Improved Sanitation*

2

Odds Ratio

$$\frac{\frac{P_1}{1-P_1}}{\frac{P_0}{1-P_0}} = 1.66$$

# Impact Evaluation: Kernel Matching Results

1

**Table:** 2006 Results for Intervention on Diarrhea for **Boys**

Variable	Treatment	(Control)	$\Delta$	S.E.
Unmatched	0.091	0.132	-0.041	(0.01)**
Matched	0.091	.151	-0.06	(0.035) <sup>†</sup>

*Note:* "Treatment" = *Improved Sanitation*

2

**Table:** 2006 Results for Intervention on Diarrhea for **Girls**

Variable	Treatment	(Control)	$\Delta$	S.E.
Unmatched	0.089	0.111	-.022	(0.01)
Matched	0.089	.1428	-0.0521	(0.03) <sup>†</sup>

*Note:* "Treatment" = *Improved Sanitation*

# Diarrhea Incidence Among Very Young Children

1

**Table: 2001 Child Diarrhea Prevalence Among  $\leq 24$  Months**

Response	Number	(%)
None	1,911	72.25
<b>Yes</b>	<b>733</b>	<b>27.7</b>
Total	2,645	100

*Source: DHS 2001*

2

**Table: 2006 Child Diarrhea Prevalence Among  $\leq 24$  Months**

Response	Number	(%)
None	1,744	81.27
<b>Yes</b>	<b>402</b>	<b>18.7</b>
Total	2,146	100

*Source: DHS 2006*

# Diarrhea Incidence Among Very Young Children

1

Table: 2006 Results for Intervention for **Children**  $\leq 24$  Months

Variable	Treatment	(Control)	$\Delta$	S.E.
Unmatched	0.151	0.203	-.052	(0.02)**
Matched	0.151	.261	-0.11	(0.05)**

*Note: "Treatment" = Improved Sanitation*

2

Odds Ratio

$$\frac{\frac{P_1}{1-P_1}}{\frac{P_0}{1-P_0}} = 1.75$$

# Nutritional Status and Diarrhea Incidence

# Impact Evaluation: Nutritional Health and Sanitation

1

**Table:** 2006 Results for **Height for Age Scores**

Variable	Treatment	(Control)	$\Delta$	S.E.
Unmatched	1884.365	1268.91	615.45	(75.44)**
Matched	1884.365	1621.09	263.27	(165.97) <sup>†</sup>

*Note:* "Treatment" = *Improved Sanitation*

2

**Table:** 2006 Results for **Weight For Age Scores**

Variable	Treatment	(Control)	$\Delta$	S.E.
Unmatched	1523.95	984.97	539	(64.78)**
Matched	1523.95	1224.52	299.42	(142.12)**

*Note:* "Treatment" = *Improved Sanitation*

# Post-Estimation: Propensity Score Distribution

# Post-Estimation: Assessing Match Quality

1

Table: Summary Statistics

Variable	Pseudo-R <sup>2</sup>	(LR $\chi^2$ )
Unmatched	0.47	2703.05
Matched	0.041	154.24

Source: DHS 2006

2

Table: Abs(Standardized Bias)

Variable	Mean	(Median)
Before Matching	28%	16%
After Matching	6.7%	2.6%

Source: DHS 2006

# Post-Estimation: Rosenbaum Bounds

I

Table: Mantel-Haenszel bounds for *Outcome = Diarrhea*

$\Gamma$	$Q_{MH+}$	$Q_{MH-}$	$p_{MH+}$	$p_{MH-}$
$\Gamma = 1$	3.05	3.05	.001	.001
$\Gamma = 1.25$	5.12	1.01	0	.15
$\Gamma = 1.50$	6.85	.53	0	.29
$\Gamma = 1.75$	8.34	1.93	0	.02
$\Gamma = 2.0$	9.66	3.16	0	0

Source: MH Bounds using STATA 10

Note:  $\Gamma = 1 \approx$  No "Hidden" Heterogeneity

Note:  $Q_{mh+}$  : Mantel-Haenszel statistic

Note:  $Q_{mh-}$  : Mantel-Haenszel statistic

Note:  $p_{mh+}$  : significance level